

Sample Questions 14

Let \mathbf{J}_n be the $n \times n$ all-ones matrix. Let \mathbf{I}_n be the $n \times n$ identity matrix.

Find S_k for $k = 0, 1, 2, 3$ and then use them to find the characteristic polynomial of \mathbf{A} .

1. Let n be a positive integer and $\omega = e^{\frac{2\pi}{n}i}$. Let $\mathbf{Q} = [\omega^{(j-1)(k-1)}]$. That is, the j, k -entry of \mathbf{Q} is $\omega^{(j-1)(k-1)}$. Show that $\frac{1}{\sqrt{n}}\mathbf{Q}$ is a unitary matrix. [This matrix is used for the Fast Fourier Transform.]

4. Let $\mathbf{A} = \mathbf{J}_n$. Find S_k for $k = 0, \dots, n$ and then use them to find the characteristic polynomial of \mathbf{J}_n .

2. Let

$$\mathbf{A} = \begin{bmatrix} 3 & 5 \\ 2 & 4 \end{bmatrix}.$$

Find S_k for $k = 0, 1, 2$ and then use them to find the characteristic polynomial of \mathbf{A} .

5. Let $\mathbf{A} = \mathbf{J}_n - \mathbf{I}_n$ be the $n \times n$ all-ones matrix. Use Problem 4 to find the characteristic polynomial of \mathbf{A} .

3. Let

$$\mathbf{A} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 4 \\ 1 & 3 & 9 \end{bmatrix}.$$

6. Let $p(x)$ be a polynomial. Let \mathbf{A} be a matrix and \mathbf{Q} an invertible matrix. Show that $\mathbf{Q}^{-1}p(\mathbf{A})\mathbf{Q} = p(\mathbf{Q}^{-1}\mathbf{A}\mathbf{Q})$.

7. Show that the Cayley–Hamilton theorem is true for diagonalizable matrices. That is, if \mathbf{A} is a diagonalizable matrix with its characteristic polynomial $p(x)$, then $p(\mathbf{A}) = \mathbf{O}$.