

國立中山大學

NATIONAL SUN YAT-SEN UNIVERSITY

線性代數 (二)

MATH 104 / GEAI 1209: Linear Algebra II

第一次期中考

March 23, 2020

Midterm 1

姓名 Name : \_\_\_\_\_

學號 Student ID # : \_\_\_\_\_

Lecturer: Jephian Lin 林晉宏

Contents: cover page,  
**6 pages** of questions,  
score page at the end

To be answered: on the test paper

Duration: **110 minutes**

Total points: **25 points** + 2 extra points

**Do not open this packet until instructed to do so.**

Instructions:

- Enter your **Name** and **Student ID #** before you start.
- Using the calculator is not allowed (and not necessary) for this exam.
- Any work necessary to arrive at an answer must be shown on the examination paper. Marks will not be given for final answers that are not supported by appropriate work.
- Clearly indicate your final answer to each question either by **underlining it or circling it**. If multiple answers are shown then no marks will be awarded.
- 可用中文或英文作答

1. Let  $\mathbf{v} = \begin{bmatrix} 5 \\ 3 \end{bmatrix}$  and  $\mathcal{E} = \{\mathbf{e}_1, \mathbf{e}_2\}$  the standard basis of  $\mathbb{R}^2$ . Let  $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2\}$  be another basis of  $\mathbb{R}^2$ , where  $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  and  $\mathbf{v}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ .

(a) [1pt] Find  $\text{Rep}_{\mathcal{E}}(\mathbf{v})$ .

(b) [1pt] Find  $\text{Rep}_{\mathcal{B}}(\mathbf{v})$ .

2. Let  $\mathbf{p} = x^2 + 2x + 3$  be a polynomial in  $\mathcal{P}_2$ , the space of all polynomials of degree at most 2.

(a) [1pt] Let  $\mathcal{B} = \{1, x, x^2\}$  be a basis of  $\mathcal{P}_2$ . Find  $\text{Rep}_{\mathcal{B}}(\mathbf{p})$ .

(b) [1pt] Let  $\mathcal{C} = \{1, x + 1, (x + 1)^2\}$  be a basis of  $\mathcal{P}_2$ . Find  $\text{Rep}_{\mathcal{C}}(\mathbf{p})$ .

(c) [1pt] Let  $\mathcal{D} = \{x^2, x, 1\}$  be a basis of  $\mathcal{P}_2$ . Find  $\text{Rep}_{\mathcal{D}}(\mathbf{p})$ .

3. Let  $\mathcal{E} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$  be the standard basis of  $\mathbb{R}^3$  and  $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  another basis of  $\mathbb{R}^3$ , where

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}.$$

- (a) [2pt] Find a matrix  $M$  such that  $M \operatorname{Rep}_{\mathcal{B}}(\mathbf{v}) = \operatorname{Rep}_{\mathcal{E}}(\mathbf{v})$  for any  $\mathbf{v} \in \mathbb{R}^3$ .

- (b) [3pt] Find a matrix  $N$  such that  $N \operatorname{Rep}_{\mathcal{E}}(\mathbf{v}) = \operatorname{Rep}_{\mathcal{B}}(\mathbf{v})$  for any  $\mathbf{v} \in \mathbb{R}^3$ .

4. Define three polynomials as follows.

$$f_1(x) = \frac{1}{2}(x-2)(x-3)$$

$$f_2(x) = -(x-1)(x-3)$$

$$f_3(x) = \frac{1}{2}(x-1)(x-2)$$

It is known that  $\mathcal{B} = \{f_1, f_2, f_3\}$  is a basis of  $\mathcal{P}_2$ , the space of all polynomials of degree at most 2.

(a) [2pt] Let  $\mathbf{p}(x) = 3x^2 + 5x + 4$ . Find  $\text{Rep}_{\mathcal{B}}(\mathbf{p})$ .

(b) [3pt] Let  $\mathcal{D} = \{1, x+1, (x+1)^2\}$  be another basis of  $\mathcal{P}_2$ . Find a matrix  $M$  such that  $M \text{Rep}_{\mathcal{D}}(\mathbf{q}) = \text{Rep}_{\mathcal{B}}(\mathbf{q})$  for any  $\mathbf{q} \in \mathcal{P}_2$ .

5. [5pt] Define a map  $f : \mathbb{R}^4 \rightarrow \mathbb{R}^3$  by  $f(\mathbf{v}) = A\mathbf{v}$ , where

$$A = \begin{bmatrix} 3 & 0 & 0 & 3 \\ 0 & 2 & 2 & 0 \\ 3 & 0 & 0 & 3 \end{bmatrix}.$$

Let  $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$  and  $\mathcal{D} = \{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$  such that

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -1 \end{bmatrix}, \mathbf{v}_4 = \begin{bmatrix} 0 \\ 1 \\ -1 \\ 0 \end{bmatrix} \text{ and.}$$

$$\mathbf{u}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \mathbf{u}_3 = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}.$$

Find  $\text{Rep}_{\mathcal{B}, \mathcal{D}}(f)$ .

6. Let  $E_{ij}$  be the  $2 \times 3$  matrix whose entries are all zeros except that the  $i, j$ -entry is one. Then

$$\mathcal{B} = \{E_{11}, E_{12}, E_{13}, E_{21}, E_{22}, E_{23}\}$$

is a basis of  $\mathcal{M}_{2 \times 3}$ , the space of all  $2 \times 3$  real matrices. Suppose  $f : \mathcal{M}_{2 \times 3} \rightarrow \mathcal{M}_{2 \times 3}$  is a homomorphism such that  $\text{Rep}_{\mathcal{B}, \mathcal{B}}(f)$  equals

$$A = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix}.$$

- (a) [1pt] Let  $M = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$ . Find  $f(M)$ .

- (b) [2pt] Find the range of  $f$ .

- (c) [2pt] Find the nullspace of  $f$ .

7. [extra 2pt] Let  $f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be a map defined by  $f(\mathbf{v}) = A\mathbf{v}$ , where

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 2 \\ 1 & 2 & 3 \end{bmatrix}.$$

Find two bases  $\mathcal{B}$  and  $\mathcal{D}$  of  $\mathbb{R}^3$  such that  $\text{Rep}_{\mathcal{B},\mathcal{D}}(f)$  is the identity matrix.

[END]

Page	Points	Score
1	5	
2	5	
3	5	
4	5	
5	5	
6	2	
Total	25 (+2)	