

國立中山大學

NATIONAL SUN YAT-SEN UNIVERSITY

線性代數 (二)

MATH 104 / GEAI 1209: Linear Algebra II

第一次期中考

March 23, 2020

Midterm 1

姓名 Name : solution

學號 Student ID # : _____

Lecturer:	Jephian Lin 林晉宏
Contents:	cover page, 6 pages of questions, score page at the end
To be answered:	on the test paper
Duration:	110 minutes
Total points:	25 points + 2 extra points

Do not open this packet until instructed to do so.

Instructions:

- Enter your **Name** and **Student ID #** before you start.
- Using the calculator is not allowed (and not necessary) for this exam.
- Any work necessary to arrive at an answer must be shown on the examination paper. Marks will not be given for final answers that are not supported by appropriate work.
- Clearly indicate your final answer to each question either by **underlining it or circling it**. If multiple answers are shown then no marks will be awarded.
- 可用中文或英文作答

1. Let $\mathbf{v} = \begin{bmatrix} 5 \\ 3 \end{bmatrix}$ and $\mathcal{E} = \{\mathbf{e}_1, \mathbf{e}_2\}$ the standard basis of \mathbb{R}^2 . Let $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2\}$ be another basis of \mathbb{R}^2 , where $\mathbf{v}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and $\mathbf{v}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$.

(a) [1pt] Find $\text{Rep}_{\mathcal{E}}(\mathbf{v})$.

$$\begin{pmatrix} 5 \\ 3 \end{pmatrix} = 5\vec{e}_1 + 3\vec{e}_2 \Rightarrow \underline{\underline{\text{Rep}_{\mathcal{E}}(\vec{v}) = \begin{pmatrix} 5 \\ 3 \end{pmatrix}}}$$

(b) [1pt] Find $\text{Rep}_{\mathcal{B}}(\mathbf{v})$.

$$\begin{pmatrix} 5 \\ 3 \end{pmatrix} = 4\vec{v}_1 + 1\vec{v}_2 \Rightarrow \underline{\underline{\text{Rep}_{\mathcal{B}}(\vec{v}) = \begin{pmatrix} 4 \\ 1 \end{pmatrix}}}$$

2. Let $\mathbf{p} = x^2 + 2x + 3$ be a polynomial in \mathcal{P}_2 , the space of all polynomials of degree at most 2.

(a) [1pt] Let $\mathcal{B} = \{1, x, x^2\}$ be a basis of \mathcal{P}_2 . Find $\text{Rep}_{\mathcal{B}}(\mathbf{p})$.

$$p = 3 \cdot 1 + 2 \cdot x + 1 \cdot x^2 \Rightarrow \underline{\underline{\text{Rep}_{\mathcal{B}}(p) = \begin{pmatrix} 3 \\ 2 \\ 1 \end{pmatrix}}}$$

(b) [1pt] Let $\mathcal{C} = \{1, x+1, (x+1)^2\}$ be a basis of \mathcal{P}_2 . Find $\text{Rep}_{\mathcal{C}}(\mathbf{p})$.

$$x^2 + 2x + 3 = a \cdot 1 + b(x+1) + c(x+1)^2$$

$$\Rightarrow a = 2, b = 0, c = 1$$

$$\Rightarrow \underline{\underline{\text{Rep}_{\mathcal{C}}(p) = \begin{pmatrix} 2 \\ 0 \\ 1 \end{pmatrix}}}$$

(c) [1pt] Let $\mathcal{D} = \{x^2, x, 1\}$ be a basis of \mathcal{P}_2 . Find $\text{Rep}_{\mathcal{D}}(\mathbf{p})$.

$$p = 1 \cdot x^2 + 2 \cdot x + 3 \cdot 1 \Rightarrow \underline{\underline{\text{Rep}_{\mathcal{D}}(p) = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}}}$$

3. Let $\mathcal{E} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$ be the standard basis of \mathbb{R}^3 and $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ another basis of \mathbb{R}^3 , where

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}.$$

- (a) [2pt] Find a matrix M such that $M \operatorname{Rep}_{\mathcal{B}}(\mathbf{v}) = \operatorname{Rep}_{\mathcal{E}}(\mathbf{v})$ for any $\mathbf{v} \in \mathbb{R}^3$.

$$M = \operatorname{Rep}_{\mathcal{B} \rightarrow \mathcal{E}}(\operatorname{id})$$

$$M = \underline{\underline{\begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix}}}$$

- (b) [3pt] Find a matrix N such that $N \operatorname{Rep}_{\mathcal{E}}(\mathbf{v}) = \operatorname{Rep}_{\mathcal{B}}(\mathbf{v})$ for any $\mathbf{v} \in \mathbb{R}^3$.

$$N = M^{-1}$$

$$(M | I) = \left(\begin{array}{ccc|ccc} 1 & -1 & 1 & 1 & 0 & 0 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right) \rightsquigarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & -1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

$$\rightsquigarrow \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 0 & 0 & 1 \end{array} \right)$$

$$\rightarrow \underline{\underline{N = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}}}$$

4. Define three polynomials as follows.

$$f_1(x) = \frac{1}{2}(x-2)(x-3)$$

$$f_2(x) = -(x-1)(x-3)$$

$$f_3(x) = \frac{1}{2}(x-1)(x-2)$$

It is known that $\mathcal{B} = \{f_1, f_2, f_3\}$ is a basis of \mathcal{P}_2 , the space of all polynomials of degree at most 2.

(a) [2pt] Let $\mathbf{p}(x) = 3x^2 + 5x + 4$. Find $\text{Rep}_{\mathcal{B}}(\mathbf{p})$.

f_1, f_2, f_3 are Lagrange polynomial based on 1, 2, 3.

$$\Rightarrow p(x) = p(1)f_1 + p(2)f_2 + p(3)f_3$$

$$p(1) = 3 + 5 + 4 = 12$$

$$p(2) = 12 + 10 + 4 = 26$$

$$p(3) = 27 + 15 + 4 = 46$$

$$\Rightarrow \text{Rep}_{\mathcal{B}}(p) = \begin{pmatrix} 12 \\ 26 \\ 46 \end{pmatrix}$$

(b) [3pt] Let $\mathcal{D} = \{1, x+1, (x+1)^2\}$ be another basis of \mathcal{P}_2 . Find a matrix M such that $M \text{Rep}_{\mathcal{D}}(\mathbf{q}) = \text{Rep}_{\mathcal{B}}(\mathbf{q})$ for any $\mathbf{q} \in \mathcal{P}_2$.

$$q(x) = q(1)f_1(x) + q(2)f_2(x) + q(3)f_3(x)$$

$$1 = 1 \cdot f_1 + 1 \cdot f_2 + 1 \cdot f_3$$

$$x+1 = 2 \cdot f_1 + 3 \cdot f_2 + 4 \cdot f_3$$

$$(x+1)^2 = 2^2 \cdot f_1 + 3^2 \cdot f_2 + 4^2 \cdot f_3$$

$$\Rightarrow \text{Rep} \Rightarrow M = \text{Rep}_{\mathcal{D}, \mathcal{B}}(\text{id}) = \begin{pmatrix} 1 & 2 & 2^2 \\ 1 & 3 & 3^2 \\ 1 & 4 & 4^2 \end{pmatrix}$$

5. [5pt] Define a map $f : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ by $f(\mathbf{v}) = A\mathbf{v}$, where

$$A = \begin{bmatrix} 3 & 0 & 0 & 3 \\ 0 & 2 & 2 & 0 \\ 3 & 0 & 0 & 3 \end{bmatrix}.$$

Let $\mathcal{B} = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4\}$ and $\mathcal{D} = \{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\}$ such that

$$\mathbf{v}_1 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \mathbf{v}_2 = \begin{bmatrix} 0 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \mathbf{v}_3 = \begin{bmatrix} 1 \\ 0 \\ 0 \\ -1 \end{bmatrix}, \mathbf{v}_4 = \begin{bmatrix} 0 \\ 1 \\ -1 \\ 0 \end{bmatrix} \text{ and.}$$

$$\mathbf{u}_1 = \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \mathbf{u}_2 = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \mathbf{u}_3 = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}.$$

Find $\text{Rep}_{\mathcal{B}, \mathcal{D}}(f)$.

$$f(\vec{v}_1) = \begin{pmatrix} 6 \\ 0 \\ 6 \end{pmatrix} = 6\vec{u}_1 + 0\vec{u}_2 + 0\vec{u}_3$$

$$f(\vec{v}_2) = \begin{pmatrix} 0 \\ 4 \\ 0 \end{pmatrix} = 0\vec{u}_1 + 4\vec{u}_2 + 0\vec{u}_3$$

$$f(\vec{v}_3) = f(\vec{v}_4) = \vec{0}$$

$$\Rightarrow \text{Rep}_{\mathcal{B}, \mathcal{D}}(f) = \begin{pmatrix} 6 & 0 & 0 & 0 \\ 0 & 4 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

6. Let E_{ij} be the 2×3 matrix whose entries are all zeros except that the i, j -entry is one. Then

$$\mathcal{B} = \{E_{11}, E_{12}, E_{13}, E_{21}, E_{22}, E_{23}\}$$

is a basis of $\mathcal{M}_{2 \times 3}$, the space of all 2×3 real matrices. Suppose $f : \mathcal{M}_{2 \times 3} \rightarrow \mathcal{M}_{2 \times 3}$ is a homomorphism such that $\text{Rep}_{\mathcal{B}, \mathcal{B}}(f)$ equals

$$A = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix}.$$

- (a) [1pt] Let $M = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$. Find $f(M)$.

$$\text{Rep}_{\mathcal{B}}(M) = \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \end{pmatrix} \quad A \cdot \text{Rep}_{\mathcal{B}}(M) = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \end{pmatrix}$$

$$\Rightarrow \underline{f(M) = \begin{pmatrix} 0 & 0 & 1 \\ 2 & 3 & 4 \end{pmatrix}}$$

- (b) [2pt] Find the range of f .
- $$\text{Colspace}(A) = \left\{ \begin{pmatrix} 0 \\ 0 \\ a \\ b \\ c \\ d \end{pmatrix} : a, b, c, d \in \mathbb{R} \right\}$$

$$\Rightarrow \underline{\text{range}(f) = \left\{ \begin{pmatrix} 0 & 0 & a \\ b & c & d \end{pmatrix} : a, b, c, d \in \mathbb{R} \right\}}$$

- (c) [2pt] Find the nullspace of f .

$$\text{Nullspace}(A) = \left\{ \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ a \\ b \end{pmatrix} : a, b \in \mathbb{R} \right\}$$

$$\Rightarrow \underline{\text{Nullspace}(f) = \left\{ \begin{pmatrix} 0 & 0 & 0 \\ 0 & a & b \end{pmatrix} : a, b \in \mathbb{R} \right\}}$$

7. [extra 2pt] Let $f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a map defined by $f(\mathbf{v}) = A\mathbf{v}$, where

$$A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 2 \\ 1 & 2 & 3 \end{bmatrix}.$$

Find two bases \mathcal{B} and \mathcal{D} of \mathbb{R}^3 such that $\text{Rep}_{\mathcal{B},\mathcal{D}}(f)$ is the identity matrix.

See ver. A.

[END]

Page	Points	Score
1	5	
2	5	
3	5	
4	5	
5	5	
6	2	
Total	25 (+2)	