

國立中山大學

NATIONAL SUN YAT-SEN UNIVERSITY

線性代數 (一)

MATH 103 / GEAI 1215: Linear Algebra I

第一次期中考

November 1, 2021

Midterm 1

姓名 Name :   solution  

學號 Student ID # : \_\_\_\_\_

Lecturer:	Jephian Lin 林晉宏
Contents:	cover page, <b>5 pages</b> of questions, score page at the end
To be answered:	on the test paper
Duration:	<b>110 minutes</b>
Total points:	<b>20 points</b> + 2 extra points

**Do not open this packet until instructed to do so.**

Instructions:

- Enter your **Name** and **Student ID #** before you start.
- Using the calculator is not allowed (and not necessary) for this exam.
- Any work necessary to arrive at an answer must be shown on the examination paper. Marks will not be given for final answers that are not supported by appropriate work.
- Clearly indicate your final answer to each question either by **underlining it or circling it**. If multiple answers are shown then no marks will be awarded.
- 可用中文或英文作答

1. Let

$$A = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 2 & 2 & 2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 2 \end{bmatrix} \text{ and } \mathbf{p} = \begin{bmatrix} 0 \\ 0 \\ 1 \\ 1 \\ 0 \end{bmatrix}.$$

(a) [1pt] Find a vector in  $\text{Row}(A)$  that is nowhere zero (每一項都不是零).

$$\underline{(1, 1, 1, 1, 2)} = (1, 1, 1, 0, 0) + (0, 0, 0, 1, 2) \in \text{Row}(A).$$

(b) [1pt] Find a vector in  $\text{Col}(A)$  that is nowhere zero.

$$\begin{pmatrix} 1 \\ 2 \\ 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \in \text{Col}(A)$$

(c) [1pt] Find a vector in  $\ker(A)$  that is nowhere zero.

$$\text{Let } \vec{x} = \begin{pmatrix} 1 \\ -1 \\ 2 \\ 2 \\ -1 \end{pmatrix}, \text{ Then } A\vec{x} = \vec{0} \Rightarrow \vec{x} \in \ker(A).$$

(d) [1pt] Find a vector in  $\mathbf{p} + \ker(A)$  that is nowhere zero.

$$\text{Pick } \vec{p} + \vec{x} = \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \\ 0 \end{pmatrix} + \begin{pmatrix} -1 \\ -1 \\ 2 \\ 2 \\ -1 \end{pmatrix} = \underline{\underline{\begin{pmatrix} -1 \\ -1 \\ 3 \\ 3 \\ -1 \end{pmatrix}}} \in \vec{p} + \ker(A)$$

(e) [1pt] Let

$$B = \begin{bmatrix} 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 2 \end{bmatrix}.$$

Find a  $3 \times 3$  matrix  $E$  such that  $EA = B$ .

$$A \xrightarrow{f_2: +(-)P_1} B \quad \text{Pick } E = \begin{pmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \text{ the elementary matrix of } f_2: +(-)P_1.$$

2. Let

$$A = \begin{bmatrix} 1 & -2 & 0 & 3 \\ 2 & -4 & 1 & 10 \\ 3 & -6 & 2 & 17 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} -4 \\ -13 \\ -22 \end{bmatrix}.$$

(a) [2pt] Find the reduced row echelon form of the augmented matrix  $[A | \mathbf{b}]$ .

$$[A | \mathbf{b}] = \left[ \begin{array}{cccc|c} 1 & -2 & 0 & 3 & -4 \\ 2 & -4 & 1 & 10 & -13 \\ 3 & -6 & 2 & 17 & -22 \end{array} \right] \longrightarrow \left[ \begin{array}{cccc|c} 1 & -2 & 0 & 3 & -4 \\ 0 & 0 & 1 & 4 & -5 \\ 0 & 0 & 2 & 8 & -10 \end{array} \right]$$

$$\longrightarrow \left[ \begin{array}{cccc|c} 1 & -2 & 0 & 3 & -4 \\ 0 & 0 & 1 & 4 & -5 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right]$$

(b) [3pt] Find  $\mathbf{p}$ ,  $\mathbf{h}_1$ ,  $\mathbf{h}_2$  such that

$$\{\mathbf{x} \in \mathbb{R}^4 : A\mathbf{x} = \mathbf{b}\} = \mathbf{p} + \text{span}(\{\mathbf{h}_1, \mathbf{h}_2\}).$$

$$\begin{cases} x_1 - 2x_2 + 3x_4 = -4 \\ x_3 + 4x_4 = -5 \end{cases}$$

令  $x_2 = x_4 = 0$  解  $A\vec{x} = \vec{b} \Rightarrow x_3 = -5, x_1 = -4 \Rightarrow \vec{p} = \begin{pmatrix} -4 \\ 0 \\ -5 \\ 0 \end{pmatrix}$

令  $x_2 = 1, x_4 = 0$  解  $A\vec{x} = \vec{0} \Rightarrow x_3 = 0, x_1 = 2 \Rightarrow \vec{h}_1 = \begin{pmatrix} 2 \\ 1 \\ 0 \\ 0 \end{pmatrix}$

令  $x_2 = 0, x_4 = 1$  解  $A\vec{x} = \vec{0} \Rightarrow x_3 = -4, x_1 = -3 \Rightarrow \vec{h}_2 = \begin{pmatrix} -3 \\ 0 \\ -4 \\ 1 \end{pmatrix}$

3. Let

$$A = \begin{bmatrix} 1 & 1 & 2 \\ -1 & -1 & -2 \\ 1 & 2 & 3 \\ -1 & -2 & -3 \end{bmatrix} \text{ and } \mathbf{b} = \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}.$$

(a) [3pt] Find  $\mathbf{w}$  and  $\mathbf{h}$  such that  $\mathbf{b} = \mathbf{w} + \mathbf{h}$  with  $\mathbf{w} \in \text{Col}(A)$  and  $\mathbf{h} \in \text{Col}(A)^\perp$ .

$$\text{Let } B = \begin{bmatrix} 1 & 1 \\ -1 & -1 \\ 1 & 2 \\ -1 & -2 \end{bmatrix} \Rightarrow \text{Col}(A) = \text{Col}(B).$$

 $\vec{w}$  = projection of  $\vec{b}$  onto  $\text{Col}(A) = \text{Col}(B)$ .

$$= B(B^T B)^{-1} B^T \vec{b}$$

$$= \begin{bmatrix} 1 & 1 \\ -1 & -1 \\ 1 & 2 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} 4 & 6 \\ 6 & 10 \end{bmatrix}^{-1} \begin{bmatrix} 1 & -1 & 1 & -1 \\ 1 & -1 & 2 & -2 \end{bmatrix} \begin{bmatrix} 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$= \begin{bmatrix} 1 & 1 \\ -1 & -1 \\ 1 & 2 \\ -1 & -2 \end{bmatrix} \frac{1}{4} \begin{bmatrix} 10 & -6 \\ -6 & 4 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 1 & 1 \\ -1 & -1 \\ 1 & 2 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} 4 \\ -2 \end{bmatrix} = \frac{1}{4} \begin{bmatrix} 2 \\ -2 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 1/2 \\ -1/2 \\ 0 \\ 0 \end{bmatrix}$$

$$\vec{h} = \vec{b} - \vec{w} = \begin{bmatrix} 1/2 \\ 1/2 \\ 0 \\ 0 \end{bmatrix}$$

(b) [2pt] Let  $\theta$  be the angle between  $\mathbf{b}$  and  $\mathbf{w}$ . Find  $\cos \theta$ .

$$\cos \theta = \frac{\langle \vec{b}, \vec{w} \rangle}{\|\vec{b}\| \cdot \|\vec{w}\|}$$

$$= \frac{1/2}{1 \cdot 1/\sqrt{2}}$$

$$= \frac{\sqrt{2}}{2}$$

4. [5pt] 數學作文：請寫一篇短文來向沒修過線性代數的朋友介紹什麼是子空間 (subspace)。

請以盡量白話的敘述、或是比喻來介紹什麼是子空間？為什麼要考慮這樣的觀念？並給一些能幫助他人理解的例子（正面的、反面的）；有必要的話可以加上一些圖來輔助說明。格式沒有限制，篇幅大約半面到一面。

(If Chinese is not your native language, you may use English or the language that you prefer.)

5. [extra 2pt] Let

$$A = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & 2 & 2 & 2 & 2 \\ 1 & 2 & 3 & 3 & 3 \\ 1 & 2 & 3 & 4 & 4 \\ 1 & 2 & 3 & 4 & 5 \end{bmatrix} \quad \text{and} \quad I = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}.$$

Find a  $5 \times 5$  matrix  $E$  such that  $E^T J E = I$ .

See ver. A

[END]

Page	Points	Score
1	5	
2	5	
3	5	
4	5	
5	2	
Total	20 (+2)	