## 線性代數 (二) MATH 104A / GEAI 1209A: Linear Algebra II

期末考 June 3, 2024 **Final Exam** 

姓名 Name :

學號 Student ID # : \_\_\_\_\_

Lecturer: Jephian Lin 林晉宏

Contents: cover page,

6 pages of questions, score page at the end

To be answered: on the test paper

Duration: 110 minutes

Total points: 20 points + 2 extra points

Do not open this packet until instructed to do so.

## **Instructions:**

- Enter your Name and Student ID # before you start.
- Using the calculator is not allowed (and not necessary) for this exam.
- Any work necessary to arrive at an answer must be shown on the examination paper. Marks will not be given for final answers that are not supported by appropriate work.
- Clearly indicate your final answer to each question either by **underlining** it or circling it. If multiple answers are shown then no marks will be awarded.
- Please answer the problems in English.

1. Let  $f: \mathbb{R}^3 \to \mathbb{R}^2$  be a linear function,  $\alpha = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  a basis of  $\mathbb{R}^3$ , and  $\beta = \{\mathbf{u}_1, \mathbf{u}_2\}$  a basis of  $\mathbb{R}^2$  such that

$$[f]_{\alpha}^{\beta} = \begin{bmatrix} 5 & 0 & 0 \\ 0 & 2 & 0 \end{bmatrix},$$

$$\mathbf{v}_{1} = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \mathbf{v}_{2} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \mathbf{v}_{3} = \begin{bmatrix} 1 \\ 4 \\ 9 \end{bmatrix}, \mathbf{u}_{1} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \text{ and } \mathbf{u}_{2} = \begin{bmatrix} 6 \\ 7 \end{bmatrix}.$$

(a) [1pt] Find  $f(\mathbf{v}_2)$ .

(b) [1pt] Find  $f(\mathbf{v}_3)$ .

(c) [1pt] Find  $f(\mathbf{v}_1 + \mathbf{v}_2)$ .

(d) [1pt] Find  $[\mathbf{v}_1 + \mathbf{v}_3]_{\alpha}$ .

(e) [1pt] Find  $f\left(\begin{bmatrix} 2\\5\\10\end{bmatrix}\right)$ .

2. Let

$$A = \begin{bmatrix} 2 & 4 \\ 4 & 2 \end{bmatrix}.$$

(a) [2pt] For the optimization problem

$$\begin{array}{ll}
\text{max} & \mathbf{x}^{\top} A \mathbf{x}, \\
\text{subject to} & \|\mathbf{x}\| = 1,
\end{array}$$

find the maximum value and the x that achieves it.

(b) [1pt] Let

$$D = \begin{bmatrix} -2 & 0 \\ 0 & 6 \end{bmatrix}.$$

Find a vector  $\mathbf{x}$  with  $\|\mathbf{x}\| = 1$  such that  $\mathbf{x}^{\top} D\mathbf{x} = 0$ .

(c) [2pt] Find a vector  $\mathbf{x}$  with  $\|\mathbf{x}\| = 1$  such that  $\mathbf{x}^{\top} A \mathbf{x} = 0$ .

3. Let

$$A = \begin{bmatrix} \frac{2}{3} & -\frac{1}{3} & -\frac{2}{3} \\ -\frac{1}{3} & \frac{2}{3} & -\frac{2}{3} \\ -\frac{2}{3} & -\frac{2}{3} & -\frac{1}{3} \end{bmatrix}.$$

(a) [4pt] Find the spectral decomposition

$$A = \sum_{j=1}^{q} \mu_j P_j.$$

(b) [1pt] Explain what does the action  $\mathbf{x} \mapsto A\mathbf{x}$  do.

4. [5pt] Mathematical essay: Write a few paragraphs to introduce *linear* regression.

Your score will be based on the following criteria.

- The definition is clear.
- Some sentences are added to explain the definition.
- Examples or pictures are included to help understanding.
- The sentences are complete.

5. [extra 5pt] Let

$$A = \begin{bmatrix} 1 & 1 & -2 & -2 \\ 1 & 1 & 2 & 2 \end{bmatrix}.$$

Find the singular value decomposition

$$A = U\Sigma V^{\top}.$$

6. [extra 2pt] Consider the  $10 \times 10$  multiplication table as a matrix

$$A = \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 9 & 10 \\ 2 & 4 & 6 & 8 & 10 & 12 & 14 & 16 & 18 & 20 \\ 3 & 6 & 9 & 12 & 15 & 18 & 21 & 24 & 27 & 30 \\ 4 & 8 & 12 & 16 & 20 & 24 & 28 & 32 & 36 & 40 \\ 5 & 10 & 15 & 20 & 25 & 30 & 35 & 40 & 45 & 50 \\ 6 & 12 & 18 & 24 & 30 & 36 & 42 & 48 & 54 & 60 \\ 7 & 14 & 21 & 28 & 35 & 42 & 49 & 56 & 63 & 70 \\ 8 & 16 & 24 & 32 & 40 & 48 & 56 & 64 & 72 & 80 \\ 9 & 18 & 27 & 36 & 45 & 54 & 63 & 72 & 81 & 90 \\ 10 & 20 & 30 & 40 & 50 & 60 & 70 & 80 & 90 & 100 \end{bmatrix}$$

Find all the eigenvalues of A.

Page	Points	Score
1	5	
2	5	
3	5	
4	5	
5	5	
6	2	
Total	20 (+7)	