

# Inverse Eigenvalue Problem of a Graph

Jephian C.-H. Lin 林晉宏


Department of Applied Mathematics, National Sun Yat-sen University

December 9, 2022  
05C50 Online Seminar, virtual

# Inverse eigenvalue problem of a graph (IEP- $G$ )

Let  $G$  be a graph. Define  $\mathcal{S}(G)$  as the family of all real symmetric matrices  $A = [a_{ij}]$  such that

$$a_{ij} \begin{cases} \neq 0 & \text{if } ij \in E(G), i \neq j; \\ = 0 & \text{if } ij \notin E(G), i \neq j; \\ \in \mathbb{R} & \text{if } i = j. \end{cases}$$




IEP- $G$ : What are the possible spectra of a matrix in  $\mathcal{S}(G)$ ?


# Inverse eigenvalue problem of a graph (IEP- $G$ )

Let  $G$  be a graph. Define  $\mathcal{S}(G)$  as the family of all real symmetric matrices  $A = [a_{ij}]$  such that

$$a_{ij} \begin{cases} \neq 0 & \text{if } ij \in E(G), i \neq j; \\ = 0 & \text{if } ij \notin E(G), i \neq j; \\ \in \mathbb{R} & \text{if } i = j. \end{cases}$$




IEP- $G$ : What are the possible spectra of a matrix in  $\mathcal{S}(G)$ ?



$$\text{spec}(A) = \{\lambda_1^{(m_1)}, \dots, \lambda_q^{(m_q)}\} \implies \mathbf{m}(A) = (m_1, \dots, m_q),$$

$$q(A) = q$$



## Questions

What are possible  $\mathbf{m}(A)$  and what are

$$M(G) = \max\{\text{mult}_\lambda(A) : A \in \mathcal{S}(G), \lambda \in \text{spec}(A)\},$$

$$q(G) = \min\{q(A) : A \in \mathcal{S}(G)\}?$$

$$\begin{bmatrix} ? & * & & & \\ * & ? & * & & \\ & * & \ddots & \ddots & \\ & & \ddots & & * \\ & & & * & ? \end{bmatrix} \in \mathcal{S}(P_n) \quad \lambda_1 < \cdots < \lambda_n$$

- $\{\text{Rows } 2 \sim n\}$  and  $\{\text{Rows } 1 \sim n-1\}$  are always independent.
- $\text{mult}(\lambda) = \text{null}(A - \lambda I) \leq 1$  for any  $A \in \mathcal{S}(P_n)$  and  $\lambda \in \mathbb{R}$ .

$$\begin{bmatrix} ? & * & & & \\ * & ? & * & & \\ & * & \ddots & \ddots & \\ & & \ddots & & * \\ & & & * & ? \end{bmatrix} \in \mathcal{S}(P_n) \quad \lambda_1 < \cdots < \lambda_n$$

Theorem (Gray and Wilson 1976; and Hald 1976)

For any set  $\Lambda = \{\lambda_1, \dots, \lambda_n\}$  of  $n$  *distinct* real numbers, there is a matrix  $A \in \mathcal{S}(P_n)$  such that  $\text{spec}(A) = \Lambda$ .

$$\begin{bmatrix} ? & * & & & * \\ * & ? & * & & \\ & * & \ddots & \ddots & \\ & & \ddots & & * \\ * & & & * & ? \end{bmatrix} \in \mathcal{S}(C_n)$$

$$\lambda_1 \leq \lambda_2 < \lambda_3 \leq \lambda_4 < \dots, \text{ or}$$

$$\lambda_1 < \lambda_2 \leq \lambda_3 < \lambda_4 \leq \lambda_5 < \dots$$

For example,  $(2, 1, 2)$  is not possible for  $C_5$ .

- $\{\text{Rows } 2 \sim n-1\}$  is always independent.
- $\text{mult}(\lambda) = \text{null}(A - \lambda I) \leq 2$  for any  $A \in \mathcal{S}(C_n)$  and  $\lambda \in \mathbb{R}$ .



$$\begin{bmatrix} ? & * & & & * \\ * & ? & * & & \\ & * & \ddots & \ddots & \\ & & \ddots & & * \\ * & & & * & ? \end{bmatrix} \in \mathcal{S}(C_n)$$

$$\lambda_1 \leq \lambda_2 < \lambda_3 \leq \lambda_4 < \cdots, \text{ or}$$

$$\lambda_1 < \lambda_2 \leq \lambda_3 < \lambda_4 \leq \lambda_5 < \cdots$$

For example,  $(2, 1, 2)$  is not possible for  $C_5$ .

### Theorem (Ferguson 1980)

*For any set  $\Lambda = \{\lambda_1, \dots, \lambda_n\}$  of  $n$  real numbers satisfying one of the conditions above, there is a matrix  $A \in \mathcal{S}(C_n)$  such that  $\text{spec}(A) = \Lambda$ .*

## Signature similarity

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 2 & 0 & 0 \\ 2 & 3 & -4 & 0 \\ 0 & -4 & 5 & -6 \\ 0 & 0 & -6 & 7 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

### Definition

A **signature matrix** is a matrix whose diagonal entries are 1 or  $-1$ . Two matrices  $A$  and  $B$  are **signature similar** if  $B = DAD$  for some signature matrix.

### Observation

Every matrix  $A \in \mathcal{S}(P_n)$  is signature similar to a matrix  $A' \in \mathcal{S}(P_n)$  whose off-diagonal entries are nonnegative.

## What do we know about the eigenvectors?

$$\begin{bmatrix} 1 & 2 & 0 & 0 \\ 2 & 3 & 4 & 0 \\ 0 & 4 & 5 & 6 \\ 0 & 0 & 6 & 7 \end{bmatrix} = QDQ^T \text{ with } Q = \begin{bmatrix} 0.33 & 0.86 & 0.38 & 0.05 \\ -0.58 & -0.13 & 0.75 & 0.28 \\ 0.63 & -0.36 & 0.17 & 0.67 \\ -0.4 & 0.34 & -0.5 & 0.69 \end{bmatrix}$$

### Theorem (Ferguson 1980)

Let  $A \in \mathcal{S}(P_n)$  with nonnegative off-diagonal entries. Suppose  $\lambda_1 < \dots < \lambda_n$  are the eigenvalues of  $A$  and  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  the corresponding orthonormal eigenbasis. Then  $(\mathbf{v}_i)_1(\mathbf{v}_i)_n$  is sign alternating for  $i = 1, \dots, n$ .

# Adjugate

Let  $A$  be an  $n \times n$  matrix.

- $A(i, j)$  is the submatrix of  $A$  by removing the  $i$ -th row and the  $j$ -th column.
- The  $i, j$ -cofactor of  $A$  is

$$c_{i,j}(A) = (-1)^{i+j} \det(A(i, j)).$$

- The **adjugate** of  $A$  is  $A^{\text{adj}} = [c_{i,j}]^T$ .
- It is known that  $AA^{\text{adj}} = A^{\text{adj}}A = \det(A)I$ , so

$$A^{\text{adj}} = \begin{cases} \det(A)A^{-1} & \text{if } \text{rank}(A) = n, \\ O & \text{if } \text{rank}(A) \leq n - 2. \end{cases}$$

# Eigenvector-eigenvalue identity

## Theorem (Eigenvector-eigenvalue identity)

Let  $A$  be an  $n \times n$  real symmetric matrix. Suppose  $\lambda_1 \leq \dots \leq \lambda_n$  are the eigenvalues of  $A$  and  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  the corresponding orthonormal eigenbasis. If  $\text{mult}(\lambda_i) = 1$ , then

$$(A - \lambda_i I)^{\text{adj}} = \left( \prod_{j \neq i} (\lambda_j - \lambda_i) \right) \mathbf{v}_i \mathbf{v}_i^\top.$$

## Corollary

When  $A \in \mathcal{S}(P_n)$  is a matrix with nonnegative off-diagonal entries,

$$\text{sgn}((\mathbf{v}_i)_1 (\mathbf{v}_i)_n) = (-1)^{1+n} \det((A - \lambda_i)(n, 1)) \prod_{j \neq i} (\lambda_j - \lambda_i).$$

## Spectrum of $A \in \mathcal{S}(C_n)$

### Theorem (Ferguson 1980)

Let  $A \in \mathcal{S}(C_n)$  with eigenvalues  $\lambda_1 \leq \dots \leq \lambda_n$ . If  $\lambda_i = \lambda_{i+1}$  and  $\lambda_j = \lambda_{j+1}$  for some  $i < j$ , then  $j - i$  is even.

### Sketch of the proof

By signature similarity, We may assume

$$A = \begin{bmatrix} A(n) & \mathbf{b} \\ \mathbf{b}^\top & c \end{bmatrix} \text{ with } \mathbf{b} = \begin{bmatrix} x \\ \mathbf{0} \\ y \end{bmatrix}$$

such that  $A(n) \in \mathcal{S}(P_{n-1})$  with **nonnegative off-diagonal** entries and eigenvalues  $\mu_1 < \dots < \mu_{n-1}$ .

(Let  $\{\mathbf{v}_1, \dots, \mathbf{v}_{n-1}\}$  be the corresponding orthonormal eigenbasis of  $A(n)$ .)

# Spectrum of $A \in \mathcal{S}(C_n)$

## Theorem (Ferguson 1980)

Let  $A \in \mathcal{S}(C_n)$  with eigenvalues  $\lambda_1 \leq \dots \leq \lambda_n$ . If  $\lambda_i = \lambda_{i+1}$  and  $\lambda_j = \lambda_{j+1}$  for some  $i < j$ , then  $j - i$  is even.

## Sketch of the proof

By the Cauchy interlacing theorem,

$$\lambda_i \leq \mu_i \leq \lambda_{i+1} = \lambda_i$$

implies  $\lambda_i = \mu_i = \lambda_{i+1}$  and similarly  $\lambda_j = \mu_j = \lambda_{j+1}$ .

# Spectrum of $A \in \mathcal{S}(C_n)$

## Theorem (Ferguson 1980)

Let  $A \in \mathcal{S}(C_n)$  with eigenvalues  $\lambda_1 \leq \dots \leq \lambda_n$ . If  $\lambda_i = \lambda_{i+1}$  and  $\lambda_j = \lambda_{j+1}$  for some  $i < j$ , then  $j - i$  is even.

## Sketch of the proof

$$A = \begin{bmatrix} A(n) & \mathbf{b} \\ \mathbf{b}^\top & c \end{bmatrix} \text{ with } \mathbf{b} = \begin{bmatrix} x \\ \mathbf{0} \\ y \end{bmatrix}$$

- Since  $\text{mult}_A(\lambda_i) = 2$  and  $\text{mult}_{A(n)}(\lambda_i) = 1$ ,  $\mathbf{b} \in \text{Col}(A(n) - \mu_i I)$ .
- $\mathbf{b} \perp \mathbf{v}_i \implies \text{sgn}(xy) = -\text{sgn}((\mathbf{v}_i)_1)(\mathbf{v}_i)_{n-1}$  (same for  $j$ )
- Since  $\text{sgn}(xy)$  is fixed and  $\text{sgn}((\mathbf{v}_j)_1)(\mathbf{v}_j)_{n-1})$  is **sign alternating**,  $j - i$  is even.



# For the IEP- $G$ , we need . . .

- Combinatorial tools: zero forcing, unique shortest path, variants of zero forcing . . .
- Theory of symmetric matrices: Cauchy interlacing theorem, eigenvector-eigenvalue identity, Rayleigh quotient, Parter–Wiener theorem, Godsil’s lemma, . . .
- Analytic tools: implicit function theorem, inverse function theorem, . . .


## An introductory article

S. M. Fallat, L. Hogben, J. C.-H. Lin, and B. Shader.

The inverse eigenvalue problem of a graph, zero forcing, and related parameters.

*Notices Amer. Math. Soc.*, 67:257–261, February, 2020.


## A book



# Inverse eigenvalue problem of a graph (IEP- $G$ )

Let  $G$  be a graph. Define  $\mathcal{S}(G)$  as the family of all real symmetric matrices  $A = [a_{ij}]$  such that

$$a_{ij} \begin{cases} \neq 0 & \text{if } ij \in E(G), i \neq j; \\ = 0 & \text{if } ij \notin E(G), i \neq j; \\ \in \mathbb{R} & \text{if } i = j. \end{cases}$$




IEP- $G$ : What are the possible spectra of a matrix in  $\mathcal{S}(G)$ ?

# Inverse eigenvalue problem of a graph (IEP- $G$ )

Let  $G$  be a graph. Define  $\mathcal{S}(G)$  as the family of all real symmetric matrices  $A = [a_{ij}]$  such that

$$a_{ij} \begin{cases} \neq 0 & \text{if } ij \in E(G), i \neq j; \\ = 0 & \text{if } ij \notin E(G), i \neq j; \\ \in \mathbb{R} & \text{if } i = j. \end{cases}$$



IEP- $G$ : What are the possible spectra of a matrix in  $\mathcal{S}(G)$ ?

# Supergraph lemma

## Lemma (BFHHLS 2017)

Let  $G$  and  $H'$  be two graphs with  $V(G) = V(H')$  and  $E(G) \subseteq E(H')$ . If  $A \in \mathcal{S}(G)$  has the **SSP**, then there is a matrix  $A' \in \mathcal{S}(H')$  such that

- $\text{spec}(A') = \text{spec}(A)$ ,
- $A'$  has the SSP, and
- $\|A' - A\|$  can be chosen arbitrarily small.


$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 3 & 0 \\ 0 & 0 & 0 & 4 \end{bmatrix} \longrightarrow \begin{bmatrix} \sim 1 & \epsilon & 0 & 0 \\ \epsilon & \sim 2 & \epsilon & 0 \\ 0 & \epsilon & \sim 3 & \epsilon \\ 0 & 0 & \epsilon & \sim 4 \end{bmatrix}$$

SSP will be defined later

## Inverse function theorem in $\mathbb{R}^2$

Fix a point  $\mathbf{a} \in \mathbb{R}^2$ . Combine two perturbations:

$$\begin{cases} \mathbf{a} + b\mathbf{e}_1 & (\text{same } y) \\ R_\theta \mathbf{a} & (\text{same norm}) \end{cases} \implies F(b, \theta) = R_\theta \mathbf{a} + b\mathbf{e}_1$$



$\frac{dF}{db, \theta}$  invertible  $\implies$  any nearby  $\mathbf{w}$  can be written as  $\mathbf{w} = F(b', \theta')$

For whatever  $y$  value nearby, there is  $\mathbf{a}'$  with  $\|\mathbf{a}'\| = \|\mathbf{a}\|$ .

# Inverse function theorem

## Theorem (Inverse function theorem)

Let  $F : U \rightarrow W$  be a smooth function. If  $\dot{F}$  at a point  $\mathbf{u}_0 \in U$  is invertible, then  $F$  is locally invertible around  $\mathbf{u}_0$ .


## Theorem (FHLS 2022)

Let  $F : U \rightarrow W$  be a smooth function. If  $\dot{F}$  at a point  $\mathbf{u}_0 \in U$  is surjective, then  $F$  is locally surjective around  $\mathbf{u}_0$ .

# Inverse function theorem in $\text{Sym}_n(\mathbb{R})$

Fix a point  $A \in \mathcal{S}(G)$ . Combine two perturbations:

$$\begin{cases} A + B & \text{(same pattern)} \\ e^{-K} A e^K & \text{(same spectrum)} \end{cases} \implies F(B, K) = e^{-K} A e^K + B$$



$\dot{F}$  surjective  $\implies$  any nearby  $M$  can be written as  $M = F(B', K')$

For whatever pattern nearby, there is  $A'$  with  $\text{spec}(A') = \text{spec}(A)$ .

## Pattern perturbation

$$F(B, K) = e^{-K} A e^K + B$$

Define  $\mathcal{S}^{\text{cl}}(G)$  as the topological closure of  $\mathcal{S}(G)$ :

$$\mathcal{S}^{\text{cl}}(G) = \{A = [a_{i,j}] \in \text{Sym}_n(\mathbb{R}) : a_{i,j} = 0 \iff \{i, j\} \in E(\overline{G})\}.$$

Let  $A \in \mathcal{S}(G)$ . Then  $A + B \in \mathcal{S}(G)$  when  $\|B\|$  is small enough.

The tangent space of  $F(B, K)$  at  $(O, O)$  with respect to  $B$  is  $\mathcal{S}^{\text{cl}}(G)$ .



# Isospectral perturbation

$$F(B, K) = e^{-K} A e^K + B$$

The function  $e^K$  is a bijection between

$\{\text{skew-symmetric matrices nearby } O\} \rightarrow \{\text{orthogonal matrices nearby } I\}$

for real matrices.

The tangent space of  $F(B, Q)$  at  $(O, O)$  with respect to  $Q$  is

$$\{-KA + AK : K \in \text{Skew}_n(\mathbb{R})\}.$$

# Strong spectral property, transversality, and surjectivity

## Definition


Let  $A$  be a real symmetric matrix. Then  $A$  has the **strong spectral property (SSP)** if  $X = O$  is the only real symmetric matrix that satisfies

$$A \circ X = I \circ X = [A, X] = O.$$

Let  $F(B, K) = e^{-K} A e^K + B$ . Then the following are equivalent:

- 1  $A$  has the SSP.
- 2  $\mathcal{S}^{\text{cl}}(G) + \{-KA + AK : K \in \text{Skew}_n(\mathbb{R})\} = \text{Sym}_n(\mathbb{R})$ .
- 3 The derivative  $\dot{F}$  is surjective.

# Illustration of the supergraph lemma



For whatever pattern nearby, there is  $A'$  with  $\text{spec}(A') = \text{spec}(A)$ .

# They must be true, right?

Let  $A \in \mathcal{S}(G)$  with the **SSP**. People *believed* that ...

- For any set of real numbers  $\Lambda'$  **nearby**  $\text{spec}(A)$ , there is a matrix  $A' \in \mathcal{S}(G)$  with  $\text{spec}(A') = \Lambda'$ .
- For any **refinement**  $\mathbf{m}'$  of  $\mathbf{m}(A)$ , there is a matrix  $A' \in \mathcal{S}(G)$  with  $\mathbf{m}(A') = \mathbf{m}'$ .
- For any  $k > q(A)$ , there is a matrix  $A' \in \mathcal{S}(G)$  with  $q(A') = k$ .

Let  $A \in \mathcal{Q}(P)$  be a nilpotent matrix with the **nSSP**. People *knew* that ...


- For any set of complex numbers  $\Lambda'$  (**invariant under conjugation**) **nearby**  $\{0, \dots, 0\}$ , there is a matrix  $A' \in \mathcal{Q}(P)$  with  $\text{spec}(A') = \Lambda'$ .

nSSP = the condition of the nilpotent-centralizer method

# Bifurcation lemma

## Theorem (FHLS 2022)

Let  $A \in \mathcal{S}(G)$  with the SSP. Then for any set of real numbers  $\Lambda'$  nearby  $\text{spec}(A)$ , there is a matrix  $A'$  with  $\text{spec}(A') = \Lambda'$ .



$$F(B, K) = e^{-K}(A + B)e^K$$

# The nSSP

## Definition

Let  $A$  be a real matrix. Then  $A$  has the **non-symmetric strong spectral property (nSSP)** if  $X = O$  is the only real matrix that satisfies

$$A \circ X = [A, X^T] = O.$$

Let  $Q^v(P)$  be the set of matrices with the same zero entries as  $P$ .

Let  $F(B, Q) = Q^{-1}AQ + B$ , where  $B \in Q^v(P)$ . Then the following are equivalent:


- 1  $A$  has the nSSP.
- 2  $Q^v(P) + \{-LA + AL : L \in \text{Mat}_n(\mathbb{R})\} = \text{Mat}_n(\mathbb{R})$ .
- 3 The derivative  $\dot{F}$  is surjective.

# Bifurcation lemma for non-symmetric matrices

## Theorem (FHLS 2022)

Let  $A \in Q(P)$  with the  $n$ SSP for some sign pattern  $P$ . Then for any set of complex numbers  $\Lambda'$  (invariant under conjugation) nearby  $\text{spec}(A)$ , there is a matrix  $A' \in Q(P)$  with  $\text{spec}(A') = \Lambda'$ .

Supergraph lemma:  $F_1(B, K) = e^{-K} A e^K + B$




Bifurcation lemma:  $F_2(B, K) = e^{-K} (A + B) e^K$

Thanks!






Supergraph lemma:  $F_1(B, K) = e^{-K} A e^K + B$



Bifurcation lemma:  $F_2(B, K) = e^{-K} (A + B) e^K$

Thanks!

# References I

-  W. Barrett, S. M. Fallat, H. T. Hall, L. Hogben, J. C.-H. Lin, and B. Shader.  
Generalizations of the Strong Arnold Property and the minimum number of distinct eigenvalues of a graph.  
*Electron. J. Combin.*, 24:#P2.40, 2017.
-  S. M. Fallat, L. Hogben, J. C.-H. Lin, and B. Shader.  
The inverse eigenvalue problem of a graph, zero forcing, and related parameters.  
*Notices Amer. Math. Soc.*, 67:257–261, February, 2020.
-  W. E. Ferguson, Jr.  
The construction of Jacobi and periodic Jacobi matrices with prescribed spectra.  
*Math. Comp.*, 35:1203–1220, 1980.

## References II



L. J. Gray and D. G. Wilson.

Construction of a Jacobi matrix from spectral data.

*Linear Algebra Appl.*, 14:131–134, 1976.



O. H. Hald.

Inverse eigenvalue problems for Jacobi matrices.

*Linear Algebra Appl.*, 14:63–85, 1976.



L. Hogben, J. C.-H. Lin, and B. Shader.

*Inverse Problems and Zero Forcing for Graphs.*

American Mathematical Society, Providence, 2022.